

FEMALE NEW PATIENT PACKAGE

The contents of this package are your first step to restore your vitality.

Please take time to read this carefully and answer all the questions as completely as possible.

Thank you for your interest in BioTE Medical®. In order to determine if you are a candidate for bio-identical testosterone pellets, we need laboratory and your history forms. We will evaluate your information prior to your consultation to determine if BioTE Medical® can help you live a healthier life. Please complete the following tasks before your appointment:

2 weeks or more before your scheduled consultation: Get your blood labs drawn at any CPL, Quest or LabCorp location. IF YOU ARE NOT INSURED OR HAVE A HIGH DEDUCTIBLE, CALL OUR OFFICE FOR SELF-PAY BLOOD DRAWS. We request the tests listed below. It is your responsibility to find out if your insurance company will cover the cost, and which lab to go to. Please note that it can take up to two weeks for your lab results to be received by our office. Please faast for 12 hours prior to blood draw.

Your blood work panel MUST include the following tests:

Estradiol **FSH** Testosterone Total TSH _T4, Total _T3, Free _T.P.O. Thyroid Peroxidase CBC _Complete Metabolic Panel _Vitamin D, 25-Hydroxy (Optional) _Vitamin B12 (Optional) _Lipid Panel (Optional) (Must be a fasting blood draw to be accurate) Female Post Insertion Labs Needed at 4, 6 or 8 Weeks based on your practitioner's choice: **FSH** _Testosterone Total Lipid Panel (Optional) (Must be a fasting blood draw to be accurate) _TSH, T4 Total, Free T3, TPO (Needed only if you've been prescribed thyroid medication) _Estradiol





FEMALE TREATMENT PLAN

- The following medications or supplements are recommended in addition to your pellet therapy.
- Please refer to the supplement brochure to help you understand why these are beneficial.
- Unless specified, these can be taken any time of day without regards to meals.

Supplements: These may be purchased in our office. When you run out they can be mailed to you for your convenience.

your convenience.		
ADK 5,000 (vitamins A, D3	and K2)	
1 a day2 a day	y forweeks, then one a day	
ADK 10,000 (vitamins A, D3	3 and K2)	
1 a day2 a day	y forweeks, then one a day	
Probiotic Take 1 a day for or	ne week, then take 2 a day starting	g week 2
Omega 3 Take 1 -4 softge	ls a daily with meal	
BioTE lodine Plus 12.5 mg	daily with food or as directed	by physician
DIM Takea day		
Whole Food Multi Take_	a day	
Meriva 500SF Take 2 a da	y MethylGua i	rd Plus Take 3 a day
Prescriptions: These have been	called into your preferred pha	rmacy
Progesterone/Prometriu	m nightly	
100mg 200	mg	
Please do not skip doses of this	s medication as it can result in v	aginal bleeding or an
increased risk for endometrial c	ancer.	
Nature-Throid/WP Thyro	id mg twice a day - upor	n waking and at 1pm. This should
be taken on an empty stomach	. Please wait 30 minutes before	e putting anything else on your
stomach. This includes coffee, fo	ood, medications, vitamins or s	supplements.
Wean off Synthroid/levot	hyroxine: alternate your desicc	ated thyroid (Nature-throid)
every other day with Synthroid/	levothyroxine for 3 weeks then	go to every day on your
desiccated thyroid.		
Spironolactone 100 mg d	aily	(other)
Wean off your antidepres	sant (see wean protocol)	_ (other)
Please call or email for any qu	estions about these recommenda	tions.
	ed a copy and understand the instruc	
Print Name	Signature	Today's Date





Symptom Checklist for Women

Name	Date:			
E-Mail:				
Symptom (please check mark)	Never	Mild	Moderate	Severe
Depressive mood Memory loss Mental confusion Decreased sex drive/libido Sleep problems Mood changes/Irritability Tension Migraine/severe headaches Difficult to climax sexually Bloating Weight gain Breast tenderness Vaginal dryness Hot flashes Night sweats				
Dry and wrinkled skin Hair falling out Cold all the time Swelling all over the body Joint pain Family History				
			NO	YES
Heart Disease Diabetes Osteoporosis Alzheimer's Disease Breast Cancer				
				,





FEMALE PATIENT QUESTIONNAIRE & HISTORY

Name:			Ioday's Dat	e:
(Last)	(First)	(Mid	dle)	
Date of Birth:	Age:	Weight:	Occupation:	
Home Address:	*			
City:			State:	Zip:
Home Phone:	Cell	Phone:	Work:	
E-Mail Address:		Ma	y we contact you via E-Mail	?()YES()NO
In Case of Emergency	Contact:		Relationship:	
Home Phone:	Cell	Phone:	Work:	
Primary Care Physiciar	n's Name:		Phone:	
Address: (Address)	(City)	(State) dow () Living with Partner	(Zip)
permission to speak to are giving us permissio	your spouse or signi on to speak with your	ficant other abou spouse or signific	ovided above, we would like t your treatment. By giving t ant other about your treatn	he information below you nent.
Spouse's Name:			Relationship:	
Home Phone:	Cell	Phone:	Work:	
Social: () I am sexually active () I want to be sexually active () I have completed () My sex has suffer () I haven't been ab	ally active. I my family. ed.	m.		
Habits: () I smoke cigarettes () I drink alcoholic b () I drink more than () I use caffeine	peverages 10 alcoholic bever	per we	eek.	







MEDICAL HISTORY

Any known drug allergies:	
Have you ever had any issues with anesthesia? () Y	es () No
If yes, please explain:	
Medications Currently Taking:	
Past Hormone Replacement Therapy:	
Nutritional/Vitamin Supplements:	
Surgeries, list all and when:	is a second of the second of t
Last menstrual period (estimate year if unknown):	
Other Pertinent Information:	
Preventative Medical Care:	Medical Illnesses:
() Medical/GYN exam in the last year.	() Polycystic Ovary Syndrome(PCOS)
() Mammogram in the last 12 months.	() High blood pressure.
() Bone density in the last 12 months.	() Heart bypass.
() Pelvic ultrasound in the last 12 months.	() High cholesterol.
High Risk Past Medical/Surgical History:	() Hypertension.
() Breast cancer.	() Heart disease.
() Uterine cancer.	() Stroke and/or heart attack.
() Ovarian cancer.	() Blood clot and/or a pulmonary emboli.
() Hysterectomy with removal of ovaries.	() Arrhythmia.
() Hysterectomy only.	() Any form of Hepatitis or HIV.
() Oophorectomy removal of ovaries.	() Lupus or other auto immune disease.
Birth Control Method:	() Fibromyalgia.
() Menopause.	() Trouble passing urine or take Flomax or Avodart.
() Hysterectomy.	() Chronic liver disease (hepatitis, fatty liver, cirrhosis).
() Tubal ligation.	() Diabetes.
() Birth control pills.	() Thyroid disease.
() Vasectomy.	() Arthritis.
() Other:	() Depression/anxiety.
	() Psychiatric disorder.
	() Cancer (type):
	Year:





Fema	le Testostero	ne and/or E	strac	liol Pellet	Insertion Co	onsent Fo	rm
Name:	webs	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,			Today's Date:		
(Las	t) (I	First)	(Mic	ldle)			
menopause. Esti mones have the	rmone pellets are hor rogen and testosteror same effects on your ions (ups and downs)	ne were made in yo body as your own	ur ovari estroger	es and adrenal g	and prior to mend	pause. Bio-ide	ntical hor-
The pellet methor	rmone pellets are pla od of hormone replac s. You will have simila	ement has been us	ed in Eu	rope and Canad	a for many years a	ind by select OI	B/GYNs in
Patients who are ment therapy. T	e pre-menopausal are estosterone is catego	advised to continury X (will cause birt	ie reliab h defect	le birth control v s) and cannot be	vhile participating e given to pregnan	in pellet hormo t women.	ne replace-
My birth con Abstinence	trol method is: (pleas Birth control pill	e circle) Hysterectomy	IUD	Menopause	Tubal ligation	Vasectomy	Other
that I may experi ated to tradition	REATMENT: I consent ence any of the comp al testosterone and/ I in the list of overall	olications to this pro or estrogen replace	ocedure	as described be	low. These side ef	fects are similar	r to those re-
sexuality (overace three weeks (est (estrogen only); exposed to testo	g, swelling, infection tive Libido); lack of e rogen pellets only); in increased growth of e esterone during their g ent (which is reversib	ffect (from lack of a ncrease in hair grow estrogen dependen gestation; growth o	absorption th on the ttumors fliver tu	on); breast tende le face, similar to li (endometrial ca mors, if already i	erness and swellin pre-menopausal ncer, breast cance present; change in	g especially in t patterns; water er); birth defect voice (which is	he first retention s in babies reversible);

and can cause bleeding. Testosterone therapy may increase one's hemoglobin and hematocrit, or thicken one's blood. This problem can be diagnosed with a blood test. Thus, a complete blood count (Hemoglobin & Hematocrit) should be done at least annually. This condition can be reversed simply by donating blood periodically.

BENEFITS OF TESTOSTERONE PELLETS INCLUDE: Increased libido, energy, and sense of well-being; increased muscle mass and strength and stamina; decreased frequency and severity of migraine headaches; decrease in mood swings, anxiety and irritability; decreased weight; decrease in risk or severity of diabetes; decreased risk of heart disease; decreased risk of Alzheimer's and dementia.

I have read and understand the above. I have been encouraged and have had the opportunity to ask any questions regarding pellet therapy. All of my questions have been answered to my satisfaction. I further acknowledge that there may be risks of testosterone and or estrogen therapy that we do not yet know, at this time, and that the risks and benefits of this treatment have been explained to me and I have been informed that I may experience complications, including one or more of those listed above. I accept these risks and benefits, and I consent to the insertion of hormone pellets under my skin. This consent is ongoing for this and all future pellet insertions.

I understand that payment is due in full at the time of service. I also understand that it is my responsibility to submit a claim to my insurance company for possible reimbursement. I have been advised that most insurance companies do not consider pellet therapy to be a covered benefit and my insurance company may not reimburse me, depending on my coverage. I acknowledge that my provider has no contracts with any insurance company and is not contractually obligated to pre-certify treatment with my insurance company or answer letters of appeal.

Print Name	Signature	Today's Date





Commonly Asked Questions

Q. What is BioTE®?

A. BioTE® is a Bio-Identical form of hormone therapy that seeks to return the hormone balance to youthful levels in men and women.

Q. How do I know if I'm a candidate for pellets?

A. Symptoms may vary widely from depression and anxiety to night sweats and sleeplessness for example. You will be given a lab slip to have blood work done which will determine your hormone levels. Once the doctor reviews and determines you are a candidate we will schedule an appointment for insertion.

Q. Do I have blood work done before each Treatment?

A. No, only initially and 4-8 weeks later to set your dosing. You may have it done again if there are significant changes.

Q. What are the pellets made from?

A. They are made from wild yams and soy. Wild yams and soy have the highest concentration of hormones of any substance. There are no known allergens associated with wild yams and soy, because once the hormone is made it is no longer yam or soy.

Q. How long will the treatment last?

A. Every 3-6 months depending on the person. Everyone is different so it depends on how you feel and what the doctor determines is right for you. If you are really active, you are under a lot of stress or it is extremely hot your treatment may not last as long. Absorption rate is based on cardiac output.

Q. Is the therapy FDA approved?

A. What the pellets are made of is FDA approved and regulated, the process of making pellets is regulated by the State Pharmacy Board, and the distribution is regulated by the DEA and Respective State Pharmacy Boards. The PROCEDURE of placing pellets is NOT an FDA approved procedure. The pellets are derived from wild yams and soy, and are all natural and bio-identical. Meaning they are the exact replication of what the body makes.

Q. How are they administered?

A. Your practitioner will implant the pellets in the fat under the skin of the hip. A small incision is made in the hip. The pellets are inserted. No stitch is required.

Q. Does it matter if I'm on birth control?

A. No, the doctor can determine what your hormone needs are even if you are on birth control.

Q. Are there any side effects?

A. The majority of side effects is temporary and typically only happens on the first dose. All are very treatable. There are no serious side effects.

Q. What if I'm already on HRT of some sort like creams, patches, pills?

A. This is an easy transition. The doctor will be able to determine your needs even though you may be currently taking these other forms of HRT.

Q. What if I've had breast cancer?

A. Breast cancer survivors and/or those who have a history of breast cancer in their family may still be a candidate; however, this is to be determined by the physician. You should schedule a consultation with the Doctor.





WHAT MIGHT OCCUR AFTER A PELLET INSERTION

A significant hormonal transition will occur in the first four weeks after the insertion of your hormone pellets. Therefore, certain changes might develop that can be bothersome.

- FLUID RETENTION: Testosterone stimulates the muscle to grow and retain water, which may result in a weight change of two to five pounds. This is only temporary. This happens frequently with the first insertion, and especially during hot, humid weather conditions.
- **SWELLING OF THE HANDS & FEET:** This is common in hot and humid weather. It may be treated by drinking lots of water, reducing your salt intake, taking cider vinegar capsules daily, (found at most health and food stores) or by taking a mild diuretic, which the office can prescribe.
- UTERINE SPOTTING/BLEEDING: This may occur in the first few months after an insertion, especially if you have been prescribed progesterone and are not taking properly: i.e. missing doses, or not taking a high enough dose. Please notify the office if this occurs. Bleeding is not necessarily an indication of a significant uterine problem. More than likely, the uterus may be releasing tissue that needs to be eliminated. This tissue may have already been present in your uterus prior to getting pellets and is being released in response to the increase in hormones.
- MOOD SWINGS/IRRITABILITY: These may occur if you were quite deficient in hormones. They will disappear when enough hormones are in your system. 5HTP can be helpful for this temporary symptom and can be purchased at many health food stores.
- FACIAL BREAKOUT: Some pimples may arise if the body is very deficient in testosterone. This lasts a short period of time and can be handled with a good face cleansing routine, astringents and toner. If these solutions do not help, please call the office for suggestions and possibly prescriptions.
- HAIR LOSS: Is rare and usually occurs in patients who convert testosterone to DHT. Dosage adjustment generally reduces or eliminates the problem. Prescription medications may be necessary in rare cases.
- HAIR GROWTH: Testosterone may stimulate some growth of hair on your chin, chest, nipples and/ or lower abdomen. This tends to be hereditary. You may also have to shave your legs and arms more often. Dosage adjustment generally reduces or eliminates the problem.

I acknowledge that I have	received a copy and understand the instruc	tions on this form.
rint Name	Signature	Today's Date





POST-INSERTION INSTRUCTIONS FOR WOMEN

- Your insertion site has been covered with two layers of bandages. Remove the outer pressure bandage any time after 24 hours. It **must** be removed as soon as it gets wet. The inner layer is either waterproof foam tape or steri-strips. They should be removed in **3 days**.
- We recommend putting an ice pack on the insertion area a couple of times for about 20 minutes each time over the next 4 to 5 hours.
- Do not take tub baths or get into a hot tub or swimming pool for **3 days**. You may shower but do not scrub the site until the incision is well healed (about 7 days).
- No major exercises for the incision area for the next **3 days**, this includes running, elliptical, squats, lunges, etc.
- The sodium bicarbonate in the anesthetic may cause the site to swell for 1-3 days.
- The insertion site may be uncomfortable for up to 2 to 3 weeks. If there is itching or redness you may take Benadryl for relief, 50 mg. orally every 6 hours. Caution this can cause drowsiness!
- You may experience bruising, swelling, and/or redness of the insertion site which may last from a few days up to 2 to 3 weeks.
- You may notice some pinkish or bloody discoloration of the outer bandage. This is normal.
- If you experience bleeding from the incision, apply firm pressure for 5 minutes.
- Please call if you have any bleeding not relieved with pressure (not oozing), as this is NOT normal.
- Please call if you have any pus coming out of the insertion site, as this is NOT normal.

Reminders:

- Remember to go for your post-insertion blood work 6 weeks after the insertion.
- Most women will need re-insertions of their pellets 3-4 months after their initial insertion.
- Please call as soon as symptoms that were relieved from the pellets start to return to make an appointment for a re-insertion. The charge for the second visit will only be for the insertion and not a consultation.

Additional Instructions:				
I acknowledge that I have received a copy and understand the instructions on this form.				
Print Name	Signature	Today's Date		

